

B82

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 47/48, 38/39, 38/46, 39/395, 38/19, 31/405 // C07K 19/00, (A61K 47/48, 38:39) (A61K 47/48, 38:46) (A61K 47/48, 38:46, 38:39) (A61K 47/48, 31:405)</b>	<b>A2</b>	<b>(11) International Publication Number: WO 99/52562</b>  <b>(43) International Publication Date: 21 October 1999 (21.10.99)</b>
<b>(21) International Application Number: PCT/US99/08335</b> <b>(22) International Filing Date: 15 April 1999 (15.04.99)</b>  <b>(30) Priority Data:</b> 60/081,863                      15 April 1998 (15.04.98)                      US  <b>(71) Applicant: LEXIGEN PHARMACEUTICALS CORP.</b> [US/US]; 125 Hartwell Avenue, Lexington, MA 02421 (US).  <b>(72) Inventor: GILLIES, Stephen, D.; 159 Sunset Road, Carlisle, MA 01741 (US).</b>  <b>(74) Agent: BRODOWSKI, Michael, H.; Testa, Hurwitz &amp; Thibault, LLP, High Street Tower, 125 High Street, Boston, MA 02110 (US).</b>		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title: ENHANCEMENT OF ANTIBODY-CYTOKINE FUSION PROTEIN MEDIATED IMMUNE RESPONSES BY CO-ADMINISTRATION WITH ANGIOGENESIS INHIBITOR</b>  <b>(57) Abstract</b>  Disclosed are compositions and methods for enhancing a cytotoxic immune response directed against a preselected cell-type in a mammal. The methods and compositions rely on a combination of an antibody-cytokine immunoconjugate and an angiogenesis inhibitor. Once administered to the mammal, the immunoconjugate induces an immune response against the preselected cell-type, for example, a cancer cell which, as a result of the synergy with the angiogenesis inhibitor, is greater than the immune response induced by the immunoconjugate alone. The methods and compositions are particularly useful at killing solid tumors or virally-infected cells in a mammal.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**ENHANCEMENT OF ANTIBODY-CYTOKINE FUSION  
PROTEIN MEDIATED IMMUNE RESPONSES BY  
CO-ADMINISTRATION WITH ANGIOGENESIS INHIBITOR**

**Reference to Related Applications**

This application claims priority to, and the benefit of U.S.S.N. 60/081,863, filed April 15, 1998, the disclosure of which is incorporated by reference herein.

**Field of the Invention**

The present invention relates generally to immunoconjugates, in particular, antibody-  
5 cytokine fusion proteins useful for targeted immune therapy and general immune stimulation.  
More specifically, the present invention relates to the use of angiogenesis inhibitors to enhance an  
antibody-cytokine fusion protein mediated immune response against a preselected cell-type, for  
example, cells in a solid tumor.

**Background of the Invention**

10       Antibodies have been used for treatment of human diseases for many years, primarily to  
provide passive immunity to viral or bacterial infection. More recently, however, antibodies and  
antibody conjugates have been used as anti-tumor agents. Anti-tumor activity has been difficult  
to demonstrate in most tumor types unless the clinical setting is one of minimal residual disease  
(Reithmuller *et al.*, LANCET 94: 1177-1183), or when the tumor is accessible to antibodies in the  
15       circulation, for example, in the case of B-lymphoma (Maloney *et al.* (1994) BLOOD 84:  
2457-2466). Solid tumors are much more refractory to antibody-mediated therapeutic  
intervention than are micrometastatic foci found in minimal residual disease settings.

Earlier studies show that the treatment of tumors with antibodies *in vivo* can be enhanced  
greatly by fusing immune stimulatory cytokines to an antibody molecule. However, antibody-  
20       cytokine fusion proteins were far less effective in destroying larger, solid tumors than they were  
for disseminated metastatic foci (Xiang *et al.* (1997) CANCER RESEARCH 57: 4948-4955, and  
Lode *et al.* (1998) BLOOD 91: 1706-1715).

Therefore, there still remains a need in the art for compositions and methods employing  
such compositions for enhancing antibody-cytokine fusion protein mediated immune responses  
25       against preselected cell-types, for example, cell-types present in solid tumors.

- 2 -

Summary of the Invention

This invention is based, in part, upon the discovery that when an immunoconjugate is administered to a mammal, it is possible to create a more potent immune response against a preselected cell-type if the immunoconjugate is administered together with an angiogenesis inhibitor. In particular, it has been found that such combinations are particularly useful in mediating the immune destruction of the preselected cell-type, such as cell-types found in solid tumors and in virally-infected cells.

In one aspect, the invention provides a method of inducing a cytocidal immune response against a preselected cell-type in a mammal. The method comprises administering to the mammal (i) an immunoconjugate comprising an antibody binding site capable of binding the preselected cell-type and a cytokine capable of inducing such an immune response against the preselected cell-type, and (ii) an angiogenesis inhibitor in an amount sufficient to enhance the immune response relative to the immune response stimulated by immunoconjugate alone.

In a preferred embodiment, the preselected cell-type can be a cancer cell present, for example, in a solid tumor, more preferably in a larger, solid tumor (i.e., greater than about 100 mm<sup>3</sup>). Alternatively, the preselected cell-type can be a virally-infected cell, for example, a human immunodeficiency virus (HIV) infected cell.

In another preferred embodiment, the angiogenesis inhibitor can be administered simultaneously with the immunoconjugate. Alternatively, the angiogenesis inhibitor can be administered prior to administration of the immunoconjugate. Furthermore, it is contemplated that the immunoconjugate can be administered together with a plurality of different angiogenesis inhibitors. Alternatively, it is contemplated that the angiogenesis inhibitor can be administered together with a plurality of different immunoconjugates.

In another aspect, the invention provides a composition for inducing a cytocidal immune response against a preselected cell-type in a mammal. The composition comprises in combination: (i) an immunoconjugate comprising an antibody binding site capable of binding the preselected cell-type, and a cytokine capable of inducing such an immune response against the preselected cell-type in the mammal, and (ii) an angiogenesis inhibitor in an amount sufficient to enhance the

- 3 -

immune response induced by the immunoconjugate of the combination relative to the immune response stimulated by the immunoconjugate alone.

In a preferred embodiment, the antibody binding site of the immunoconjugate preferably comprises, an immunoglobulin heavy chain or an antigen binding fragment thereof. The  
5 immunoglobulin heavy chain preferably comprises, in an amino-terminal to carboxy-terminal direction, an immunoglobulin variable (VH) region domain capable of binding a preselected antigen, an immunoglobulin constant heavy 1 (CH1) domain, an immunoglobulin constant heavy 2 (CH2) domain, and optionally may further include an immunoglobulin constant heavy 3 (CH3) domain. In a more preferred embodiment, the immunoconjugate is a fusion protein comprising an  
10 immunoglobulin heavy chain or an antigen binding fragment thereof fused via a polypeptide bond to the cytokine. Accordingly, a preferred antibody-cytokine fusion protein comprises, in an amino-terminal to carboxy-terminal direction, (i) the antibody binding site comprising an immunoglobulin variable region capable of binding a cell surface antigen on the preselected cell-type, an immunoglobulin CH1 domain, an immunoglobulin CH2 domain (optionally a CH3  
15 domain), and (ii) the cytokine. Methods for making and using such fusion proteins are described in detail in Gillies *et al.* (1992) PROC. NATL. ACAD. SCI. USA 89: 1428-1432; Gillies *et al.* (1998) J. IMMUNOL. 160:6195-6203; and U.S. Patent No. 5,650,150.

The immunoglobulin constant region domains (i.e., the CH1, CH2 and/or CH3 domains) may be the constant region domains normally associated with the variable region domain in a  
20 naturally occurring antibody. Alternatively, one or more of the immunoglobulin constant region domains may derived from antibodies different from the antibody used as a source of the variable region domain. In other words, the immunoglobulin variable and constant region domains may be derived from different antibodies, for example, antibodies derived from different species. See, for example, U.S. Patent No. 4,816,567. Furthermore, the immunoglobulin variable regions may  
25 comprise framework region (FR) sequences derived from one species, for example, a human, and complementarity determining region (CDR) sequences interposed between the FRs, derived from a second, different species, for example, a mouse. Methods for making and using such chimeric immunoglobulin variable regions are disclosed, for example, in U.S. Patent Nos. 5,225,539 and 5,585,089.

- 4 -

The antibody-based immunoconjugates preferably further comprise an immunoglobulin light chain which preferably is covalently bonded to the immunoglobulin heavy chain by means of, for example, a disulfide bond. The variable regions of the linked immunoglobulin heavy and light chains together define a single and complete binding site for binding the preselected antigen. In  
5 other embodiments, the immunoconjugates comprise two chimeric chains, each comprising at least a portion of an immunoglobulin heavy chain fused to a cytokine. The two chimeric chains preferably are covalently linked together by, for example, one or more interchain disulfide bonds.

The invention thus provides fusion proteins in which the antigen-binding specificity and activity of an antibody is combined with the potent biological activity of a cytokine. A fusion  
10 protein of the present invention can be used to deliver the cytokine selectively to a target cell *in vivo* so that the cytokine can exert a localized biological effect in the vicinity of the target cell. In a preferred embodiment, the antibody component of the fusion protein specifically binds an antigen on a cancer cell and, as a result, the fusion protein exerts localized anti-cancer activity. In an alternative preferred embodiment, the antibody component of the fusion protein specifically  
15 binds a virus-infected cell, such as an HIV-infected cell, and, as a result, the fusion protein exerts localized anti-viral activity.

Preferred cytokines that can be incorporated into the immunoconjugates of the invention include, for example, tumor necrosis factors, interleukins, colony stimulating factors, and lymphokines. Preferred tumor necrosis factors include, for example, tissue necrosis factor  $\alpha$   
20 (TNF $\alpha$ ). Preferred interleukins include, for example, interleukin-2 (IL-2), interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-7 (IL-7), interleukin-12 (IL-12), interleukin-15 (IL-15) and interleukin-18 (IL-18). Preferred colony stimulating factors include, for example, granulocyte-macrophage colony stimulating factor (GM-CSF) and macrophage colony stimulation factor (M-CSF). Preferred lymphokines include, for example, lymphotoxin (LT). Other useful cytokines  
25 include interferons, including IFN- $\alpha$ , IFN- $\beta$  and IFN- $\gamma$ , all of which have immunological effects, as well as anti-angiogenic effects, that are independent of their anti-viral activities.

Preferred angiogenesis inhibitors useful in the practice of the invention include, for example, endostatin, angiostatin, peptides having binding affinity for  $\alpha_v\beta_3$  integrin, antibodies or fragments thereof having binding affinity for  $\alpha_v\beta_3$  integrin, peptides with binding affinity for an  
30 epidermal growth factor (EGF) receptor, antibodies or fragments thereof having binding affinity

- 5 -

for an EGF receptor, COX-2 inhibitors, fumagillin, thalidomide, anti-angiogenic cytokines, for example, IFN- $\alpha$ , IFN- $\beta$  and IFN- $\gamma$ , and a cytokine fusion protein comprising such an anti-angiogenic cytokine.

Also provided are preferred dosages and administration regimes for administering the  
5 immunoconjugates in combination with the angiogenesis inhibitors.

- 6 -

*Brief Description of the Drawings*

The foregoing and other objects, features, and advantages of the present invention, as well as the invention itself, may be more fully understood from the following description of preferred embodiments, when read together with the accompanying drawings, in which:

5        FIG. 1 is a schematic representation of an exemplary immunoconjugate useful in the practice of the invention.

FIG. 2A and 2B are graphs depicting the expression of human EpCAM in transfected mouse Lewis lung carcinoma (LLC) cells as analyzed by fluorescence-activated cell sorting (FACS). Equal numbers of transfected cells were stained either with a secondary fluorescein isothiocyanate (FITC)-labeled anti-human Fc specific antibody alone (Panel A), or first stained  
10        with a huKS-huIL2 antibody fusion protein followed by the FITC-labeled anti-human Fc specific antibody (Panel B);

FIG. 3 is a line graph depicting the effects on subcutaneous tumors of an antibody-cytokine fusion protein administered either alone or in combination with a second antibody-cytokine fusion protein in which the cytokine has anti-angiogenic activity. Treatment for 5 days  
15        was initiated 13 days after implantation of LLC cells. The mice were treated with phosphate buffered saline (open diamonds); 15 µg/day of huKS-mur2a-muIL2 fusion protein alone (closed squares); 10 µg/day of a huKS-mur2a-muIL12 fusion protein (closed triangles); and a combination of 7.5 µg/day of the huKS-mur2a-muIL2 fusion protein and 5 µg/day of the  
20        huKS-mur2a-muIL12 fusion protein (crosses);

FIG. 4 is a line graph depicting the effects on subcutaneous tumors of an antibody-cytokine fusion protein administered either alone or in combination with an endostatin fusion protein. The size of CT26/EpCAM subcutaneous tumors were monitored in mice treated with phosphate buffered saline (closed diamonds), a muFc-muEndo fusion protein (closed squares), a  
25        huKS-huγ4-huIL2 fusion protein (closed diamond), and a combination of the muFc-muEndo fusion protein and the huKS-huγ4-huIL2 fusion protein (crosses); and

FIG. 5 is a line graph depicting the effect on subcutaneous tumors of an antibody-cytokine fusion protein administered either alone or in combination with indomethacin. The size of LLC-EpCAM subcutaneous tumors were monitored in mice treated with phosphate buffered saline



- 7 -

(closed diamonds), a huKS-huγ1-huIL2 fusion protein (closed squares), indomethocin (closed triangles), and a combination of the huKS-huγ1-huIL2 fusion protein and indomethocin (crosses).

- 8 -

Detailed Description of the Invention

It has now been discovered that cytotoxic immune responses initiated by an immunoconjugate against a preselected cell-type can be significantly enhanced by administering the immunoconjugate together with an angiogenesis inhibitor. The combined therapy is particularly effective in mediating the immune destruction of a diseased tissue, such as, an established tumor or virally-infected cells. The present invention describes methods for making and using useful immunoconjugates, as well as assays useful for testing their pharmacokinetic activities in pre-clinical *in vivo* animal models when they are combined with suitable angiogenesis inhibitors.

As used herein, the term "cytotoxic immune response" is understood to mean any immune response in a mammal, either humoral or cellular in nature, that is stimulated by the immunoconjugate of the invention and which either kills or otherwise reduces the viability of a preselected cell-type in the mammal. The immune response may include one or more cell types, including T cells, natural killer (NK) cells and macrophages.

As used herein, the term "immunoconjugate" is understood to mean a conjugate of (i) an antibody binding site having binding specificity for, and capable of binding a surface antigen on a cancer cell or a virally-infected cell, and (ii) a cytokine that is capable of inducing or stimulating a cytotoxic immune response against the cancer or virally-infected cell. Accordingly, the immunoconjugate is capable of selectively delivering the cytokine to a target cell *in vivo* so that the cytokine can mediate a localized immune response against the target cell. For example, if the antibody component of the immunoconjugate selectively binds an antigen on a cancer cell, for example, a cancer cell in a solid tumor, in particular, a larger solid tumor of greater than about 100 mm<sup>3</sup>, the immunoconjugate exerts localized anti-cancer activity. Alternatively, if the antibody component of the immunoconjugate selectively binds an antigen on a virally-infected cell, such as a human immunodeficiency virus (HIV) infected cell, the immunoconjugate exerts localized anti-viral activity.

As used herein, the term "antibody binding site" is understood to mean at least a portion of an immunoglobulin heavy chain, for example, an immunoglobulin variable region capable of binding the preselected cell-type. The antibody binding site also preferably comprises at least a

- 9 -

portion of an immunoglobulin constant region including, for example, a CH1 domain, a CH2 domain, and optionally, a CH3 domain. Furthermore, the immunoglobulin heavy chain may be associated, either covalently or non-covalently, to an immunoglobulin light comprising, for example, an immunoglobulin light chain variable region and optionally light chain constant region.

5 Accordingly, it is contemplated that the antibody binding site may comprise an intact antibody or a fragment thereof capable of binding the preselected cell-type.

With regard to the immunoconjugate, it is contemplated that the antibody fragment may be linked to the cytokine by a variety of ways well known to those of ordinary skill in the art. For example, the antibody binding site preferably is linked via a polypeptide bond to the cytokine in a fusion protein construct. Alternatively, the antibody binding site may be chemically coupled to

10 the cytokine via reactive groups, for example, sulfhydryl groups, within amino acid sidechains present within the antibody binding site and the cytokine.

As used herein, the term "cytokine" is understood to mean any protein or peptide, analog or functional fragment thereof, which is capable of stimulating or inducing a cytotoxic immune response against a preselected cell-type, for example, a cancer cell or a virally-infected cell, in a mammal. Accordingly, it is contemplated that a variety of cytokines can be incorporated into the immunoconjugates of the invention. Useful cytokines include, for example, tumor necrosis factors, interleukins, lymphokines, colony stimulating factors, interferons including species variants, truncated analogs thereof which are capable of stimulating or inducing such cytotoxic

15 immune responses. Useful tumor necrosis factors include, for example, TNF  $\alpha$ . Useful lymphokines include, for example, LT. Useful colony stimulating factors include, for example, GM-CSF and M-CSF. Useful interleukins include, for example, IL-2, IL-4, IL-5, IL-7, IL-12, IL-15 and IL-18. Useful interferons, include, for example, IFN- $\alpha$ , IFN- $\beta$  and IFN- $\gamma$ .

The gene encoding a particular cytokine of interest can be cloned *de novo*, obtained from an available source, or synthesized by standard DNA synthesis from a known nucleotide

25 sequence. For example, the DNA sequence of LT is known (see, for example, Nedwin *et al.* (1985) NUCLEIC ACIDS RES. 13:6361), as are the sequences for IL-2 (see, for example, Taniguchi *et al.* (1983) NATURE 302:305-318), GM-CSF (see, for example, Gasson *et al.* (1984) SCIENCE

- 10 -

266:1339-1342), and TNF  $\alpha$  (see, for example, Nedwin *et al.* (1985) NUCLEIC ACIDS RES. 13:6361).

In a preferred embodiment, the immunoconjugates are recombinant fusion proteins produced by conventional recombinant DNA methodologies, i.e., by forming a nucleic acid  
5 construct encoding the chimeric immunoconjugate. The construction of recombinant antibody-cytokine fusion proteins has been described in the prior art. See, for example, Gillies *et al.* (1992) PROC. NATL. ACAD. SCI. USA 89: 1428-1432; Gillies *et al.* (1998) J. IMMUNOL. 160:6195-6203; and U.S. Patent No 5,650,150. Preferably, a gene construct encoding the immunoconjugate of the invention includes, in 5' to 3' orientation, a DNA segment encoding an immunoglobulin heavy  
10 chain variable region domain, a DNA segment encoding an immunoglobulin heavy chain constant region, and a DNA encoding the cytokine. The fused gene is assembled in or inserted into an expression vector for transfection into an appropriate recipient cell where the fused gene is expressed. The hybrid polypeptide chain preferably is combined with an immunoglobulin light chain such that the immunoglobulin variable region of the heavy chain ( $V_H$ ) and the  
15 immunoglobulin variable region of the light chain ( $V_L$ ) combine to produce a single and complete site for binding a preselected antigen. In a preferred embodiment, the immunoglobulin heavy and light chains are covalently coupled, for example, by means of an interchain disulfide bond. Furthermore, two immunoglobulin heavy chains, either one or both of which are fused to a cytokine, can be covalently coupled, for example, by means of one or more interchain disulfide  
20 bonds.

Figure 1 shows a schematic representation of an exemplary immunoconjugate 10. In this embodiment, cytokine molecules 2 and 4 are peptide bonded to the carboxy termini 6 and 8 of CH3 regions 10 and 12 of antibody heavy chains 14 and 16.  $V_L$  regions 26 and 28 are shown paired with  $V_H$  regions 18 and 20 in a typical IgG configuration, thereby providing two antigen  
25 binding sites 30 and 32 at the amino terminal ends of immunoconjugate 10 and two cytokine receptor-binding sites 40 and 42 at the carboxy ends of immunoconjugate 10. Of course, in their broader aspects, the immunoconjugates need not be paired as illustrated or only one of the two immunoglobulin heavy chains need be fused to a cytokine molecule.

- 11 -

Immunoconjugates of the invention may be considered chimeric by virtue of two aspects of their structure. First, the immunoconjugate is chimeric in that it includes an immunoglobulin heavy chain having antigen binding specificity linked to a given cytokine. Second, an immunoconjugate of the invention may be chimeric in the sense that it includes an

5 immunoglobulin variable region (V) and an immunoglobulin constant region (C), both of which are derived from different antibodies such that the resulting protein is a V/C chimera. For example, the variable and constant regions may be derived from naturally occurring antibody molecules isolatable from different species. See, for example, U.S. Patent 4,816,567. Also embraced are constructs in which either or both of the immunoglobulin variable regions comprise

10 framework region (FR) sequences and complementarity determining region (CDR) sequences derived from different species. Such constructs are disclosed, for example, in Jones *et al.* (1986) NATURE 321: 522-525, Verhoyen *et al.* (1988) SCIENCE 239: 1534-1535, and U.S. Patent Nos. 5,225,539 and 5,585,089. Furthermore, it is contemplated that the variable region sequences may be derived by screening libraries, for example, phage display libraries, for variable region

15 sequences that bind a preselected antigen with a desired affinity. Methods for making and screening phage display libraries are disclosed, for example, in Huse *et al.* (1989) SCIENCE 246: 1275-1281 and Kang *et al.* (1991) PROC. NATL. ACAD. SCI. USA 88: 11120-11123.

The immunoglobulin heavy chain constant region domains of the immunoconjugates can be selected from any of the five immunoglobulin classes referred to as IgA (Ig $\alpha$ ), IgD (Ig $\delta$ ), IgE (Ig $\epsilon$ ), IgG (Ig $\gamma$ ), and IgM (Ig $\mu$ ). However, immunoglobulin heavy chain constant regions from

20 the IgG class are preferred. Furthermore, it is contemplated that the immunoglobulin heavy chains may be derived from any of the IgG antibody subclasses referred to in the art as IgG1, IgG2, IgG3 and IgG4. As is known, each immunoglobulin heavy chain constant region comprises four or five domains. The domains are named sequentially as follows: CH1-hinge-CH2-CH3-(-

25 CH4). CH4 is present in IgM, which has no hinge region. The DNA sequences of the heavy chain domains have cross homology among the immunoglobulin classes, for example, the CH2 domain of IgG is homologous to the CH2 domain of IgA and IgD, and to the CH3 domain of IgM and IgE. The immunoglobulin light chains can have either a kappa ( $\kappa$ ) or lambda ( $\lambda$ ) constant chain. Sequences and sequence alignments of these immunoglobulin regions are well

30 known in the art (see, for example, Kabat *et al.*, "Sequences of Proteins of Immunological

- 12 -

*Interest*," U.S. Department of Health and Human Services, third edition 1983, fourth edition 1987, Huck *et al.* (1986) NUC. ACIDS RES. 14: 1779-1789).

In preferred embodiments, the variable region is derived from an antibody specific for a preselected cell surface antigen (an antigen associated with a diseased cell such as a cancer cell or virally-infected cell), and the constant region includes CH1, and CH2 (and optionally CH3) domains from an antibody that is the same or different from the antibody that is the source of the variable region. In the practice of this invention, the antibody portion of the immunoconjugate preferably is non-immunogenic or is weakly immunogenic in the intended recipient. Accordingly, the antibody portion, as much as possible, preferably is derived from the same species as the intended recipient. For example, if the immunoconjugate is to be administered to humans, the constant region domains preferably are of human origin. See, for example, U.S. Patent No. 4,816,567. Furthermore, when the immunoglobulin variable region is derived from a species other than the intended recipient, for example, when the variable region sequences are of murine origin and the intended recipient is a human, then the variable region preferably comprises human FR sequences with murine CDR sequences interposed between the FR sequences to produce a chimeric variable region that has binding specificity for a preselected antigen but yet while minimizing immunoreactivity in the intended host. The design and synthesis of such chimeric variable regions are disclosed in Jones *et al.* (1986) NATURE 321: 522-525, Verhoyen *et al.* (1988) SCIENCE 239: 1534-1535, and U.S. Patent Nos. 5,225,539 and 5,585,089. The cloning and expression of a humanized antibody-cytokine fusion protein, KS-1/4 anti-EpCAM antibody - IL-12 fusion protein, as well as its ability to eradicate established colon carcinoma metastases has been described in Gillies *et al.* (1998) J. IMMUNOL. 160: 6195-6203.

The gene encoding the cytokine is joined, either directly or by means of a linker, for example, by means of DNA encoding a (Gly<sub>4</sub>-Ser)<sub>3</sub> linker in frame to the 3' end of the gene encoding the immunoglobulin constant region (e.g., a CH2 or CH3 exon). In certain embodiments, the linker can comprise a nucleotide sequence encoding a proteolytic cleavage site. This site, when interposed between the immunoglobulin constant region and the cytokine, can be designed to provide for proteolytic release of the cytokine at the target site. For example, it is well known that plasmin and trypsin cleave after lysine and arginine residues at sites that are accessible to the proteases. Many other site-specific endoproteases and the amino acid sequences

- 13 -

they cleave are well-known in the art. Preferred proteolytic cleavage sites and proteolytic enzymes that are reactive with such cleavage sites are disclosed in U.S. Patent Nos. 5,541,087 and 5,726,044.

5 The nucleic acid construct optionally can include the endogenous promoter and enhancer for the variable region-encoding gene to regulate expression of the chimeric immunoglobulin chain. For example, the variable region encoding genes can be obtained as DNA fragments comprising the leader peptide, the VJ gene (functionally rearranged variable (V) regions with joining (J) segment) for the light chain, or VDJ gene for the heavy chain, and the endogenous promoter and enhancer for these genes. Alternatively, the gene encoding the variable region can  
10 be obtained apart from endogenous regulatory elements and used in an expression vector which provides these elements.

Variable region genes can be obtained by standard DNA cloning procedures from cells that produce the desired antibody. Screening of the genomic library for a specific functionally rearranged variable region can be accomplished with the use of appropriate DNA probes such as  
15 DNA segments containing the J region DNA sequence and sequences downstream. Identification and confirmation of correct clones is achieved by sequencing the cloned genes and comparison of the sequence to the corresponding sequence of the full length, properly spliced mRNA.

The target antigen can be a cell surface antigen of a tumor or cancer cell, a virus-infected cell or another diseased cell. Genes encoding appropriate variable regions can be obtained  
20 generally from immunoglobulin-producing lymphoid cell lines. For example, hybridoma cell lines producing immunoglobulin specific for tumor associated antigens or viral antigens can be produced by standard somatic cell hybridization techniques well known in the art (see, for example, U.S. Pat. No. 4,196,265). These immunoglobulin producing cell lines provide the source of variable region genes in functionally rearranged form. The variable region genes  
25 typically will be of murine origin because this murine system lends itself to the production of a wide variety of immunoglobulins of desired specificity. Furthermore, variable region sequences may be derived by screening libraries, for example, phage display libraries, for variable region sequences that bind a preselected antigen with a desired affinity. Methods for making and

- 14 -

screening phage display libraries are disclosed, for example, in Huse *et al.* (1989) SCIENCE 246: 1275-1281 and Kang *et al.* (1991) PROC. NATL. ACAD. SCI. USA 88: 11120-11123.

5 The DNA fragment encoding containing the functionally active variable region gene is linked to a DNA fragment containing the gene encoding the desired constant region (or a portion thereof). Immunoglobulin constant regions (heavy and light chain) can be obtained from antibody-producing cells by standard gene cloning techniques. Genes for the two classes of human light chains ( $\kappa$  and  $\lambda$ ) and the five classes of human heavy chains ( $\alpha$ ,  $\delta$ ,  $\epsilon$ ,  $\gamma$  and  $\mu$ ) have been cloned, and thus, constant regions of human origin are readily available from these clones.

10 The fused gene encoding the hybrid immunoglobulin heavy chain is assembled or inserted into an expression vector for incorporation into a recipient cell. The introduction of the gene construct into plasmid vectors can be accomplished by standard gene splicing procedures. The chimeric immunoglobulin heavy chain can be co-expressed in the same cell with a corresponding immunoglobulin light chain so that a complete immunoglobulin can be expressed and assembled simultaneously. For this purpose, the heavy and light chain constructs can be placed in the same  
15 or separate vectors.

Recipient cell lines are generally lymphoid cells. The preferred recipient cell is a myeloma (or hybridoma). Myelomas can synthesize, assemble, and secrete immunoglobulins encoded by transfected genes and they can glycosylate proteins. Particularly preferred recipient or host cells include Sp2/0 myeloma which normally does not produce endogenous immunoglobulin, and  
20 mouse myeloma NS/0 cells. When transfected, the cell produces only immunoglobulin encoded by the transfected gene constructs. Transfected myelomas can be grown in culture or in the peritoneum of mice where secreted immunoconjugate can be recovered from ascites fluid. Other lymphoid cells such as B lymphocytes can be used as recipient cells.

25 There are several methods for transfecting lymphoid cells with vectors containing the nucleic acid constructs encoding the chimeric immunoglobulin chain. For example, vectors may be introduced into lymphoid cells by spheroblast fusion (see, for example, Gillies *et al.* (1989) BIOTECHNOL. 7: 798-804). Other useful methods include electroporation or calcium phosphate precipitation (see, for example, Sambrook *et al.* eds (1989) "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Press).



- 15 -

Other useful methods of producing the immunoconjugates include the preparation of an RNA sequence encoding the construct and its translation in an appropriate *in vivo* or *in vitro* expression system. It is contemplated that the recombinant DNA methodologies for synthesizing genes encoding antibody-cytokine fusion proteins, for introducing the genes into host cells, for  
5 expressing the genes in the host, and for harvesting the resulting fusion protein are well known and thoroughly documented in the art. Specific protocols are described, for example, in Sambrook *et al.* eds (1989) "*Molecular Cloning: A Laboratory Manual*," Cold Spring Harbor Press.

It is understood that the chemically coupled immunoconjugates may be produced using a  
10 variety of methods well known to those skilled in the art. For example, the antibody or an antibody fragment may be chemically coupled to the cytokine using chemically reactive amino acid side chains in the antibody or antibody fragment and the cytokine. The amino acid side chains may be covalently linked, for example, via disulfide bonds, or by means of homo- or hetero-bifunctional crosslinking reagents including, for example, N-succinimidyl 3-(2-  
15 pyridyldithio)propionate, *m*-maleimidobenzoyl-N-hydroxysuccinate ester, *m*-maleimidobenzoyl-N-hydroxysulfosuccinimide ester, and 1,4-di-[3'-(2'-pyridylthio) propionamido] butane, all of which are available commercially from Pierce, Rockford, IL.

It is understood that the term "angiogenesis inhibitor" as used herein, refers to any molecule that reduces or inhibits the formation of new blood vessels in a mammal. With regard to  
20 cancer therapy, the angiogenesis inhibitor reduces or inhibits the formation of new blood vessels in or on a tumor, preferably in or on a solid tumor. It is contemplated that useful angiogenesis inhibitors may be identified using a variety of assays well known and used in the art. Such assays include, for example, the bovine capillary endothelial cell proliferation assay, the chick chorioallantoic membrane (CAM) assay or the mouse corneal assay. However, the CAM assay is  
25 preferred (see, for example, O'Reilly *et al.* (1994) CELL 79: 315-328 and O'Reilly *et al.* (1997) CELL 88: 277-285). Briefly, embryos with intact yolks are removed from fertilized three day old white eggs and placed in a petri dish. After incubation at 37°C, 3% CO<sub>2</sub> for three days, a methylcellulose disk containing the putative angiogenesis inhibitor is applied to the chorioallantoic

- 16 -

membrane of an individual embryo. After incubation for about 48 hours, the chorioallantoic membranes were observed under a microscope for evidence of zones of inhibition.

Numerous angiogenesis inhibitors are well known and thoroughly documented in the art. Examples of angiogenesis inhibitors useful in the practice of the invention include, for example, protein/peptide inhibitors of angiogenesis such as: angiostatin, a proteolytic fragment of plasminogen (O'Reilly *et al.* (1994) CELL 79: 315-328, and U.S. Patent Nos. 5,733,876; 5,837,682; and 5,885,795); endostatin, a proteolytic fragment of collagen XVIII (O'Reilly *et al.* (1997) CELL 88: 277-285 and U.S. Patent No. 5,854,205); peptides containing the RGD tripeptide sequence and capable of binding the  $\alpha_v\beta_3$  integrin (Brooks *et al.* (1994) CELL 79: 1157-1164); and certain antibodies and antigen binding fragments thereof and peptides that interact with the  $\alpha_v\beta_3$  integrin found on tumor vascular epithelial cells (Brooks *et al.*, *supra*) or the EGF receptor (Ciardello *et al.*, (1996) J. NATL. CANCER INST. 88: 1770-1776). Examples of other angiogenesis inhibitors include: COX-2 inhibitors (Masferrer *et al.* (1998) PROC. AMER. ASSOC. CANCER RES. 39: 271); fumagillin and analogues such as AGM-1470 (Ingber *et al.* (1990) NATURE 348: 555-557); and other small molecules such as thalidomide (D'Amato *et al.* (1994) PROC. NATL. ACAD. SCI. USA 91: 4082-4085). Endostatin and angiostatin, however, currently are most preferred.

Several cytokines including species variants and truncated analogs thereof have also been reported to have anti-angiogenic activity and thus are useful in the practice of the invention. Examples include IL-12, which reportedly works through an IFN- $\gamma$ -dependent mechanism (Voest *et al.* (1995) J. NATL. CANC. INST. 87: 581-586); IFN- $\gamma$  itself, which induces a chemokine (IP-10) with angiostatic activity (Arenberg *et al.* (1996) J. EXP. MED. 184: 981-992). Thus IL-12, IFN- $\gamma$  and IP-10 represent angiogenesis inhibitors at different points of the same inhibitory pathway. Other interferons, especially IFN- $\alpha$ , have been shown to be anti-angiogenic alone or in combination with other inhibitors (Brem *et al.* (1993) J. PEDIATR. SURG. 28: 1253-1257). The interferons IFN- $\alpha$ , IFN- $\beta$  and IFN- $\gamma$  all have immunological effects, as well as anti-angiogenic properties, that are independent of their anti-viral activities. Another cytokine, GM-CSF, reportedly inhibits angiogenesis through the induction of angiostatin (Kumar *et al.* (1998) PROC. AMER. ASSOC. CANCER RES. 39: 271).

- 17 -

As used herein, it is understood that an antibody portion of the immunoconjugate specifically binds a preselected antigen, a cytokine specifically binds a receptor for the cytokine, or an angiogenesis inhibitor specifically binds a receptor for the inhibitor, if the binding affinity for the antigen or receptor is greater than  $10^5 \text{ M}^{-1}$ , and more preferably greater than  $10^7 \text{ M}^{-1}$ . As used  
5 herein, the terms angiostatin, endostatin, TNF, IL, GM-CSF, M-CSF, LT, and IFN not only refer to intact proteins but also to bioactive fragments and/or analogs thereof. Bioactive fragments refer to portions of the intact protein that have at least 30%, more preferably at least 70%, and most preferably at least 90% of the biological activity of the intact proteins. Analogs refer to  
10 species and allelic variants of the intact protein, or amino acid replacements, insertions or deletions thereof that have at least 30%, more preferably at least 70%, and most preferably at least 90% of the biological activity of the intact protein.

Angiogenesis inhibitors may be co-administered simultaneously with the immunoconjugate, or administered separately by different routes of administration. Compositions of the present invention may be administered by any route which is compatible with the particular  
15 molecules. Thus, as appropriate, administration may be oral or parenteral, including intravenous and intraperitoneal routes of administration.

The compositions of the present invention may be provided to an animal by any suitable means, directly (e.g., locally, as by injection, implantation or topical administration to a tissue locus) or systemically (e.g., parenterally or orally). Where the composition is to be provided  
20 parenterally, such as by intravenous, subcutaneous, ophthalmic, intraperitoneal, intramuscular, buccal, rectal, vaginal, intraorbital, intracerebral, intracranial, intraspinal, intraventricular, intrathecal, intracisternal, intracapsular, intranasal or by aerosol administration, the composition preferably comprises part of an aqueous or physiologically compatible fluid suspension or solution. Thus, the carrier or vehicle is physiologically acceptable so that in addition to delivery  
25 of the desired composition to the patient, it does not otherwise adversely affect the patient's electrolyte and/or volume balance. The fluid medium for the agent thus can comprise normal physiologic saline (e.g., 9.85% aqueous NaCl, 0.15M, pH 7-7.4).

Preferred dosages of the immunoconjugate per administration are within the range of  $0.1 \text{ mg/m}^2$  -  $100 \text{ mg/m}^2$ , more preferably,  $1 \text{ mg/m}^2$  -  $20 \text{ mg/m}^2$ , and most preferably  $2 \text{ mg/m}^2$  -  $6 \text{ mg/m}^2$ . Preferred dosages of the angiogenesis inhibitor will depend generally upon the type of  
30

- 18 -

angiogenesis inhibitor used, however, optimal dosages may be determined using routine experimentation. Administration of the immunoconjugate and/or the angiogenesis inhibitor may be by periodic bolus injections, or by continuous intravenous or intraperitoneal administration from an external reservoir (for example, from an intravenous bag) or internal (for example, from a bioerodable implant). Furthermore, it is contemplated that the immunoconjugate of the invention may also be administered to the intended recipient together with a plurality of different angiogenesis inhibitors. It is contemplated, however, that the optimal combination of immunoconjugates and angiogenesis inhibitors, modes of administration, dosages may be determined by routine experimentation well within the level of skill in the art.

10       A variety of methods can be employed to assess the efficacy of combined therapy using antibody-cytokine fusion proteins and angiogenesis inhibitors on immune responses. For example, the animal model described in Example 1, or other suitable animal model, can be used by a skilled artisan to test which angiogenesis inhibitors, or combinations of angiogenesis inhibitors, are most effective in acting synergistically with an immunoconjugate, for example, an antibody-cytokine fusion protein (for example, an antibody-IL2 fusion protein) to enhance the immune destruction of established tumors. The angiogenesis inhibitor, or combination of angiogenesis inhibitors, can be administered prior to, or simultaneously with, the course of immunoconjugate therapy and the effect on the tumor can be conveniently monitored by volumetric measurement. Further, as novel angiogenesis inhibitors are identified, a skilled artisan will be able to use the methods described  
15       herein to assess the potential of these novel inhibitors to enhance the anti-cancer activity of antibody-cytokine fusion proteins.  
20

      Alternatively, following therapy, tumors can be excised, sectioned and stained via standard histological methods, or via specific immuno-histological reagents in order to assess the effect of the combined therapy on immune response. For example, simple staining with hematoxylin and eosin can reveal differences in lymphocytic infiltration into the solid tumors which is indicative of a cellular immune response. Furthermore, immunostaining of sections with antibodies to specific classes of immune cells can reveal the nature of an induced response. For example, antibodies that bind to CD45 (a general leukocyte marker), CD4 and CD8 (for T cell subclass identification), and NK1.1 (a marker on NK cells) can be used to assess the type of immune response that has  
25       been mediated by the immunoconjugates of the invention.  
30

- 19 -

Alternatively, the type of immune response mediated by the immunoconjugates can be assessed by conventional cell subset depletion studies described, for example, in Lode *et al.* (1998) BLOOD 91: 1706-1715. Examples of depleting antibodies include those that react with T cell markers CD4 and CD8, as well as those that bind the NK markers NK1.1 and asialo GM.

- 5 Briefly, these antibodies are injected to the mammal prior to initiating antibody-cytokine treatment at fairly high doses (for example, at a dose of about 0.5 mg/mouse), and are given at weekly intervals thereafter until the completion of the experiment. This technique can identify the cell-types necessary to elicit the observed immune response in the mammal.

- 10 In another approach, the cytotoxic activity of splenocytes isolated from animals having been treated with the combination therapy can be compared with those from the other treatment groups. Splenocyte cultures are prepared by mechanical mincing of recovered, sterile spleens by standard techniques found in most immunology laboratory manuals. See, for example, Coligan *et al.* (eds) (1988) "Current Protocols in Immunology," John Wiley & Sons, Inc. The resulting cells then are cultured in a suitable cell culture medium (for example, DMEM from GIBCO) containing
- 15 serum, antibiotics and a low concentration of IL-2 (~10 U/mL). For example, in order to compare NK activity, 3 days of culture normally is optimal, whereas, in order to compare T cell cytotoxic activity, 5 days of culture normally is optimal. Cytotoxic activity can be measured by radioactively labeling tumor target cells (for example, LLC cells) with <sup>51</sup>Cr for 30 min. Following removal of excess radiolabel, the labeled cells are mixed with varying concentrations of cultured
- 20 spleen cells for 4 hr. At the end of the incubation, the <sup>51</sup>Cr released from the cells is measured by a gamma counter which is then used to quantitate the extent of cell lysis induced by the immune cells. Traditional cytotoxic T lymphocyte (or CTL) activity is measured in this way.

The invention is illustrated further by the following non-limiting examples.

#### **Example 1. Animal Model.**

- 25 A murine cancer model was developed to study the effect of combining antibody-cytokine fusion proteins and angiogenesis inhibitors in mediating effective immune responses against a tumor. The antibody-cytokine fusion proteins used in the following examples bind EpCAM, a human tumor antigen found on most epithelial derived tumors. (see, Perez and Walker (1989) J. IMMUNOL. 142: 3662-3667). In order to test the efficacy in an immuno-competent murine model,
- 30 it was necessary to express the human antigen on the surface of a mouse tumor cell that is

- 20 -

syngeneic with the mouse host. Lewis lung carcinoma (LLC) cells, a well known mouse lung cancer cell line, was selected for this purpose. As a result, the human tumor antigen, EpCAM, was expressed on the surface of LLC cells.

LLC cells were transfected with an expression plasmid containing a cDNA encoding  
5 human EpCAM antigen (recognized by the KS-1/4 antibody as described in Vurki *et al.* (1984) CANCER RES. 44:681), and driven by the cytomegalovirus (CMV) early promoter (Immunogen, Carlsbad, CA). The KS antigen (KSA or EpCAM) was cloned by PCR from cDNA prepared from the human prostate carcinoma cells, LnCAP. The forward primer had the oligonucleotide sequence 5' TCTAGAGCAGCATGGCGCCCCCGC (SEQ ID NO: 1), in which the ATG is the  
10 translation initiation codon, and the reverse primer had the oligonucleotide sequence 5' CTCGAGTTATGCATTGAGTTCCCT (SEQ ID NO: 2), where TTA is the anti-codon of the translation termination. The EpCAM cDNA was cloned into a retroviral vector pLNCS (Clontech, Palo Alto, CA) and transfection performed according to established protocols (Ausubel *et al.*, (eds) "Current Protocols in Molecular Biology", John Wiley & Sons). Briefly,  
15 the packaging cell line PA317 (ATCC CRL 9078) was transfected with pLNCX-EpCAM by calcium phosphate co-precipitation, and the conditioned medium containing the virus used to transfect LLC cells. G418 (Sigma Chemical Co.) was added to the transfected cells at 1 mg/mL to select for stable clones. Clones expressing human EpCAM antigen (LLC-Ep) were identified by immunostaining and fluorescence activated cell sorting (FACS) analysis.

20 As depicted in FIG. 1, LLC-Ep clones stained first with a hu-KS-IL2 antibody fusion protein (see Example 2 below) followed by a fluorescein isothiocyanate (FITC)-labeled anti-human Fc specific antibody (Jackson ImmunoResearch Laboratories, West Grove, PA), exhibited a fairly uniform level of expression of human EpCAM. The level of expression in these clones was well above the level observed with clones stained with the FITC-labeled anti-human Fc  
25 specific antibody alone.

In order to show that expression of a human cell-surface protein did not increase the immunogenicity of LLC-Ep cells, C57Bl/6 mice were injected subcutaneously with varying numbers of cells. All mice were found to develop rapidly progressive tumors after injection with  $5 \times 10^5$  cells, with roughly the same growth kinetics observed with the parental LLC cell line. All  
30 animals became moribund and were sacrificed to avoid unnecessary suffering.

- 21 -

**Example 2. Preparation of Antibody-cytokine or Antibody-angiogenesis Inhibitor Fusion Proteins.**

A variety of antibody-cytokine fusion proteins are discussed in the following examples. In particular, Example 3 discloses the use of humanized KS-murine  $\gamma$ 2a-murine IL2 (huKS-mu $\gamma$ 2a - muIL2) and humanized KS-murine  $\gamma$ 2a -murine IL12 (huKS-mu $\gamma$ 2a -muIL12) fusion proteins. Example 4 discloses the use of the huKS-mu $\gamma$ 2a -muIL2 fusion protein together with murine Fc-murine angiostatin (muFc-muAngio) and murine Fc-murine endostatin (muFc - muEndo) fusion proteins. Example 5 discloses the use of a humanized KS-human  $\gamma$ 4-human IL2 (huKS-hu $\gamma$ 4-huIL2) and muFc-muEndo fusion proteins. Finally, Example 6 discloses the use of humanized KS-human hu $\gamma$ 1-human IL2 (huKS-hu $\gamma$ 1-huIL2) fusion protein with indomethacin. The construction of these fusion proteins is discussed below.

**huKS-hu $\gamma$ 1-huIL2**

A gene encoding huKS-hu $\gamma$ 1-huIL2 fusion protein was prepared and expressed essentially as described in Gillies *et al.* (1998) J. IMMUNOL. 160:6195-6203 and U.S. Patent No. 5,650,150. Briefly, humanized variable regions of the mouse KS-1/4 antibody (Varki *et al.*, (1984) CANCER RES. 44:681-687) were modeled using the methods disclosed in Jones *et al.* (1986) NATURE 321: 522-525, which involved the insertion of the CDRs of each KS 1/4 variable region into the consensus framework sequences of the human variable regions with the highest degree of homology. Molecular modeling with a Silicon Graphics Indigo work station implementing BioSym software confirmed that the shapes of the CDRs were maintained. The protein sequences then were reverse translated, and genes constructed by the ligation of overlapping oligonucleotides.

The resulting variable regions were inserted into an expression vector containing the constant regions of the human  $\kappa$  light chain and the human C $\gamma$ 1 heavy chain essentially as described in Gillies *et al.* (1992) PROC. NATL. ACAD. SCI. USA 89:1428-1432, except that the metallothionein promoters and immunoglobulin heavy chain enhancers were replaced by the CMV promoter/enhancer for the expression of both chains. Fusions of the mature sequences of IL-2 to the carboxy terminus of the human heavy chains were prepared as described in Gillies *et al.*

- 22 -

(1992) PROC. NATL. ACAD. SCI. USA 89:1428-1432, except that the 3' untranslated regions of the IL-2 gene was derived from the SV40 poly(A) region.

The IL-2 fusion protein was expressed by transfection of the resulting plasmid into NS/0 myeloma cell line with selection medium containing 0.1  $\mu$ M methotrexate. Briefly, in order to  
5 obtain stably transfected clones, plasmid DNA was introduced into the mouse myeloma NS/0 cells by electroporation. NS/0 cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum. About  $5 \times 10^6$  cells were washed once with PBS and resuspended in 0.5 mL PBS. Ten  $\mu$ g of linearized plasmid DNA then was incubated with the cells in a Gene  
10 Pulser Cuvette (0.4 cm electrode gap, BioRad) on ice for 10 min. Electroporation was performed using a Gene Pulser (BioRad, Hercules, CA) with settings at 0.25 V and 500  $\mu$ F. Cells were allowed to recover for 10 min. on ice, after which they were resuspended in growth medium and then plated onto two 96 well plates. Stably transfected clones were selected by growth in the presence of 100 nM methotrexate (MTX), which was introduced two days post-transfection. The  
15 cells were fed every 3 days for three more times, and MTX-resistant clones appeared in 2 to 3 weeks.

Expressing clones were identified by Fc or cytokine ELISA using the appropriate antibodies (see, for example, Gillies *et al.* (1989) BIOTECHNOL. 7: 798-804). The resulting fusion protein was purified by binding, and elution from protein A Sepharose (Pharmacia), in accordance with the manufacturer's instructions.

20 **huKS-huy4-huIL2**

A gene encoding the huKS.huy4.huIL2 fusion protein was constructed and expressed essentially as described in U.S.S.N. 09/256,156, filed February 24, 1999, which claims priority to U.S.S.N. 60/075,887, filed February 25, 1998.

Briefly, an Igy4 version of the huKS-huy1-huIL2 fusion protein, described above, was  
25 prepared by removing the immunoglobulin constant region Cy1 gene fragment from the huKS-huy1-huIL2 expression vector and replacing it with the corresponding sequence from the human Cy4 gene. Sequences and sequence alignments of the human heavy chain constant regions Cy1, Cy2, Cy3, and Cy4 are disclosed in Huck *et al.* (1986) NUC. ACIDS RES. 14: 1779-1789.



- 23 -

The swapping of the Cy1 and Cy4 fragments was accomplished by digesting the original Cy1-containing plasmid DNA with Hind III and Xho I and purifying a large 7.8 kb fragment by agarose gel electrophoresis. A second plasmid DNA containing the Cy4 gene was digested with Hind III and Nsi I and a 1.75 kb fragment was purified. A third plasmid containing the human IL-2 cDNA and SV40 poly A site, fused to the carboxyl terminus of the human Cy1 gene, was digested with Xho I and Nsi I and the small 470 bp fragment was purified. All three fragments were ligated together in roughly equal molar amounts and the ligation product was used to transform competent *E. coli*. The ligation product was used to transform competent *E. coli* and colonies were selected by growth on plates containing ampicillin. Correctly assembled recombinant plasmids were identified by restriction analyses of plasmid DNA preparations from isolated transformants and digestion with Fsp I was used to discriminate between the Cy1 (no Fsp I) and Cy4 (one site) gene inserts.

The final vector, containing the Cy4-IL2 heavy chain replacement, was introduced into NS/0 mouse myeloma cells by electroporation (0.25 V and 500  $\mu$ F) and transfectants were selected by growth in medium containing methotrexate (0.1  $\mu$ M). Cell clones expressing high levels of the huKS-hu $\gamma$ 4-huIL2 fusion protein were identified, expanded, and the fusion protein purified from culture supernatants using protein A Sepharose chromatography. The purity and integrity of the Cy4 fusion protein was determined by SDS-polyacrylamide gel electrophoresis. IL-2 activity was measured in a T-cell proliferation assay (Gillis *et al.* (1978) J. IMMUNOL. 120:2027-2032) and was found to be identical to that of the  $\gamma$ 1-construct.

#### **huKS-mu $\gamma$ 2a-muIL2**

A gene encoding the huKS-mu $\gamma$ 2a-muIL2 fusion protein was constructed by replacing the human antibody constant regions and human IL-2 of the huKS-hu $\gamma$ 1-huIL2 fusion protein, as described above, with the corresponding murine sequences. Specifically, the human Cy1-IL2 DNA was replaced with a murine Cy2a cDNA fragment fused to a DNA encoding murine IL-2. Briefly, the VH region of the huKS was joined in frame to the murine  $\gamma$ 2a cDNA by performing overlapping PCR using overlapping oligonucleotide primers:

(sense) 5' CC GTC TCC TCA GCC AAA ACA ACA GCC CCA TCG GTC (SEQ ID NO: 3);

- 24 -

(antisense) 5' *GG GGC TGT TGT TTT GGC* TGA GGA GAC GGT GAC TGA CG (SEQ ID NO: 4);

(sense) 5' C TTA AGC CAG ATC CAG TTG GTG CAG (SEQ ID NO: 5); and

(antisense) 5' CC CGG GGT CCG GGA GAA GCT CTT AGT C (SEQ ID NO: 6).

5 The oligonucleotides of SEQ ID NOS: 3 and 4 were designed to hybridize to the junction of the V<sub>H</sub> domain of huKS and the constant region of murine  $\gamma$ 2a cDNA (in italics). In the first round of PCR, there were two separate reactions. In one reaction, the V<sub>H</sub> of huKS DNA was used as the template with the oligonucleotides of SEQ ID NOS: 4 and 5. The primer of SEQ ID NO: 5 introduced an AflII (CTTAAG) restriction site upstream of the sequence encoding the mature  
10 amino terminus of huKS V<sub>H</sub> (in bold). In another reaction, murine  $\gamma$ 2a cDNA was used as the template with the oligonucleotides SEQ ID NOS: 3 and 6. The primer of SEQ ID NO: 6 hybridized to the cDNA encoding the region around the C-terminus of  $\gamma$ 2a and introduced a XmaI (CCCGGG) restriction site for subsequent ligation to the muIL2 cDNA. PCR products from the two reactions were mixed and subjected to a second round of PCR, using the oligonucleotides of  
15 SEQ ID NOS: 5 and 6. The resulting PCR product was cloned, and upon sequence verification, the AflII-XmaI fragment encoding the V<sub>H</sub> of huKS and the murine  $\gamma$ 2a constant region was used for ligation to the DNA encoding the signal peptide at the AflII site and the muIL2 cDNA at the XmaI site.

20 The murine IL2 cDNA was cloned from mRNA of murine peripheral blood mononuclear cells using the oligonucleotides set forth in SEQ ID NOS: 7 and 8, namely:

(sense) 5' GGC CCG GGT AAA **GCA CCC** ACT TCA AGC TCC (SEQ ID NO. 7); and

(antisense) 5' CCCTCGAGTTATTGAGGGCTTGTTG (SEQ ID NO. 8).

The primer of SEQ ID NO: 7 adapted the muIL2 (sequence in bold) to be joined to mu  $\gamma$ 2a at the XmaI restriction site (CCCGGG). The primer of SEQ ID NO: 8 introduced an XhoI  
25 restriction site (CTCGAG) immediately after the translation termination codon (antisense in bold).

Similarly, the variable light (V<sub>L</sub>) domain of huKS was joined to the mu  $\kappa$  cDNA sequence by overlapping PCR. The overlapping oligonucleotides used included

(sense) 5' G GAA ATA AAA CGG GCT GAT GCT GCA CCA ACT G (SEQ ID NO. 9);

- 25 -

(antisense) 5' *GC AGC ATC AGC CCGTT TTA TTT CCA GCT TGG TCC* (SEQ ID NO. 10);

(sense) 5' C TTA AGC GAG ATC GTG CTG ACC CAG (SEQ ID NO. 11); and

(antisense) 5' CTC GAG CTA ACA CTC ATT CCT GTT GAA GC (SEQ ID NO. 12).

The oligonucleotides were designed to hybridize to the junction of the V<sub>L</sub> of huKS and the  
5 constant region of murine  $\kappa$  cDNA (in italics). In the first round of PCR, there were two separate  
reactions. In one reaction, the V<sub>L</sub> of huKS DNA was used as template, with the oligonucleotides  
set forth in SEQ ID NOS. 10 and 11, which introduced an AflII (CTTAAG) restriction site  
upstream of the sequence encoding the mature amino terminus of huKS V<sub>L</sub> (in bold). In the other  
reaction, murine  $\kappa$  cDNA was used as template, with the oligonucleotides set forth in SEQ ID  
10 NOS. 9 and 12, which introduced an XhoI restriction site after the translation termination codon  
(antisense in bold).

PCR products from the two reactions were mixed and subjected to a second round of  
PCR using the oligonucleotide primers set forth in SEQ ID NOS. 11 and 12. The resultant PCR  
product was cloned, and upon sequence verification, the AflII-XhoI fragment encoding the V<sub>L</sub> of  
15 huKS and the murine  $\kappa$  constant region was ligated to the DNA encoding the signal peptide at the  
AflII site.

Both the murine heavy and light chain sequences were used to replace the human  
sequences in pdHL7. The resulting antibody expression vector, containing a dhfr selectable  
marker gene, was electroporated (0.25V, 500 $\mu$ F) into murine NS/O myeloma cells and clones  
20 selected by culturing in medium containing 0.1  $\mu$ M methotrexate. Transfected clones, resistant to  
methotrexate, were tested for secretion of antibody determinants by standard ELISA methods.  
The fusion proteins were purified via protein A Sepharose chromatography according to the  
manufacturers instructions.

#### huKS-muy2a-muIL12

25 A gene encoding the huKS-muy2a-muIL12 fusion protein was constructed and expressed  
essentially as described in U.S.S.N. 08/986,997, filed December 8, 1997, and Gillies *et al.* (1998)  
J. IMMUNOL. 160:6195-6203. Briefly, this was accomplished by fusing the murine p35 IL-12  
subunit cDNA to the huKS-muy2a heavy chain coding region prepared previously. The resulting  
vector then was transfected into an NS/O myeloma cell line pre-transfected with, and capable of

- 26 -

expressing p40 IL-12 subunit. In other words, a cell line was transfected with p40 alone and a stable, high expressing cell was selected, which was then used as a recipient for transfection by the p35 containing fusion protein (i.e., sequential transfection).

The murine p35 and p40 IL-12 subunits were isolated by PCR from mRNA prepared from spleen cells activated with Concanavalin A (5 µg/mL in culture medium for 3 days). The PCR primers used to isolate the p35 encoding nucleic acid sequence which also adapted the p35 cDNA as an XmaI-XhoI restriction fragment included:

5' CCCCAGGTAGGGTCATTCCAGTCTCTGG (SEQ ID NO: 13); and

5' CTCGAGTCAGGCGGAGCTCAGATAGC (SEQ ID NO: 14).

The PCR primer used to isolate the p40 encoding nucleic acid sequence included:

5' TCTAGACCATGTGTCCTCAGAAGCTAAC (SEQ ID NO: 15); and

5' CTCGAGCTAGGATCGGACCCTGCAG (SEQ ID NO: 16).

A plasmid vector (pdHL7-huKS-mur2a-p35) was constructed as described (Gillies *et al.* J. IMMUNOL. METHODS 125:191) that contained a dhfr selectable marker gene, a transcription unit encoding a humanized KS antibody light chain, and a transcription unit encoding a murine heavy chain fused to the p35 subunit of mouse IL-12. The fusion was achieved by ligation of the XmaI to XhoI fragment of the adapted p35 subunit cDNA, to a unique XmaI site at the end of the CH3 exon of the murine  $\gamma$ 2a gene prepared previously. Both the H and L chain transcription units included a cytomegalovirus (CMV) promoter (in place of the metallothionein promoter in the original reference) at the 5' end and, a polyadenylation site at the 3' end.

A similar vector (pNC-p40) was constructed for expression of the free p40 subunit which included a selectable marker gene (neomycin resistant gene) but still used the CMV promoter for transcription. The coding region in this case included the natural leader sequence of the p40 subunit for proper trafficking to the endoplasmic reticulum and assembly with the fusion protein. Plasmid pNC-p40 was electroporated into cells, and cells were plated and selected in G418-containing medium. In this case, culture supernatants from drug-resistant clones were tested by ELISA for production of p40 subunit.

- 27 -

The pdHL7-huKS-muy2a-p35 expression vector was electroporated into the NS/0 cell line already expressing murine p40, as described in Gillies *et al.* (1998) J. IMMUNOL. 160: 6195-6203. Transfected clones resistant to methotrexate were tested for secretion of antibody determinants and mouse IL-12 by standard ELISA methods. The resulting protein was purified by binding to, and elution from a protein A Sepharose column in accordance with the manufacturers instructions.

#### muFc-muEndo

A gene encoding the muFc-muEndo fusion protein was constructed and expressed essentially as described in U.S.S.N. 60/097,883, filed August 25, 1998.

10 Briefly, murine endostatin and murine Fc were expressed as a muFc-muEndostatin fusion protein. PCR was used to adapt the endostatin gene for expression in the pdCs-muFc(D<sub>4</sub>K) vector (Lo *et al.* (1998) PROTEIN ENGINEERING 11:495-500) which contains an enterokinase recognition site Asp4-Lys (LaVallie *et al.* (1993) J. BIOL. CHEM. 268: 23311-23317). The forward primer was 5'-C CCC AAG CTT CAT ACT CAT CAG GAC TTT C (SEQ ID NO: 15 17), where the AAGCTT (*Hind*III site) was followed by sequence (in bold) encoding the N-terminus of endostatin. The reverse primer was 5'-CCC CTC GAG CTA TTT GGA GAA AGA GGT C (SEQ ID NO: 18), which was designed to put a translation STOP codon (anticodon, CTA) immediately after the C-terminus of endostatin, and this was followed by an *Xho*I site (CTCGAG).

20 The PCR product was cloned and sequenced, and the *Hind*III-*Xho*I fragment encoding endostatin was ligated into the pdCs-muFc(D<sub>4</sub>K) vector. Stable NS/0 clones expressing muFc(D<sub>4</sub>K)-muEndo were selected and assayed using an anti-muFc ELISA. The resulting fusion protein was expressed and purified via Protein A Sepharose chromatography.

#### muFc-muAngio

25 A gene encoding the muFc-muAngio fusion protein was constructed and expressed essentially as described in U.S.S.N. 60/097,883, filed August 25, 1998.

Briefly, murine angiostatin and murine Fc were expressed as a muFc-muAngiostatin fusion protein. PCR was used to adapt the angiostatin sequence NA, kindly provided by the laboratory

- 28 -

of Dr. Judah Folkman, Childrens Hospital, Boston, MA, for expression in the pdCs-Fc(D<sub>4</sub>K) vector (Lo *et al.* (1998) PROTEIN ENGINEERING 11:495-500). The forward primer was 5'-C  
 CCC AAG CTT GTG TAT CTG TCA GAA TGT AAG CCC TCC TGT CTC TGA GCA  
 (SEQ ID NO: 19), where the AAGCTT (*Hind*III site) was followed by sequence (in bold)  
 5 encoding the N-terminus of angiostatin. The reverse primer was 5'-CCC CTC GAG CTA CCC  
 TCC TGT CTC TGA GCA (SEQ ID NO: 20), which was designed to put a translation STOP  
 codon (anticodon, CTA) immediately after the C-terminus of angiostatin, and this was followed  
 by an *Xho*I site (CTCGAG).

The PCR product was cloned and sequenced, and the *Hind*III-*Xho*I fragment encoding  
 10 angiostatin was ligated into the pdCs-muFc(D<sub>4</sub>K) vector. Stable NS/0 clones expressing  
 muFc(D<sub>4</sub>K)-muAngio were selected and assayed using an anti-muFc ELISA. The resulting fusion  
 protein was expressed and purified by Protein A Sepharose chromatography.

**Example 3. Combination Therapy Using KS-IL2 and KS-IL12 fusion proteins For The Treatment of LLC-Ep Tumors.**

15 Female C57Bl/6 mice were injected subcutaneously in the mid-back with LLC-Ep cells (5  
 x 10<sup>5</sup> per mouse) grown in cell culture. After about two weeks, animals with palpable tumors in  
 the range of 150-400 mm<sup>3</sup> were divided into four groups, with an equal distribution of tumor sizes  
 between the groups. The animals were treated as follows: in group 1, animals received PBS only  
 (control group); in group 2, animals received only the huKS-muγ2a-muIL2 fusion protein; in  
 20 group 3, animals received only the huKS-muγ2a-muIL12 fusion protein; and in group 4, the  
 animals received both the huKS-muγ2a-muIL2 and the huKS-muγ2a-muIL12 fusion proteins.  
 Tumor growth was monitored by volumetric measurements until animals in the control group  
 became moribund and were euthanized. Tumor volumes were measured with calipers and  
 calculated as  $V = 4 \pi/3 (0.5L \times 0.5W \times 0.5H)$ , where L is the length, W is the width, and H is the  
 25 height of the tumor.

The results are summarized in FIG. 3. Mice treated with PBS are represented by open  
 diamonds, mice treated with 15 μg/day of huKS-muγ2a-muIL2 fusion protein are represented by  
 closed squares, mice treated with 10 μg/day of a huKS-muγ2a-muIL12 fusion protein are  
 represented by closed triangles, and mice treated with a combination of 7.5 μg/day of the

- 29 -

huKS-muγ2a-muIL2 fusion protein and 5 μg/day of the huKS-muγ2a-muIL12 fusion protein are represented by crosses.

As illustrated in FIG. 3, treatment with the huKS-muγ2a-muIL2 fusion protein (15 μg for 5 consecutive days) did not delay or reduce the growth of LLC-Ep tumors (closed squares). Only a slight anti-tumor effect was seen in mice treated with the huKS-muγ2a-muIL12 fusion protein alone at 10 μg per dose for five consecutive days (closed triangles). However, when the two fusion proteins were combined using half of the original amounts for each (7.5 μg of huKS-muγ2a-muIL2 and 5 μg of huKS-muγ2a-muIL12, respectively), a striking growth delay was observed (crosses) suggesting a synergistic effect between the two fusion proteins.

**Example 4. Therapy Using An Antibody-cytokine Fusion Protein And A Combination Of Angiostatin And Endostatin.**

Female C57Bl/6 mice can be injected subcutaneously in the mid-back with LLC-Ep cells ( $5 \times 10^5$  per mouse) grown in cell culture. After about two weeks, animals with palpable tumors in the range of 150-400 mm<sup>3</sup> are divided into four groups and treated as follows: (1) animals receiving only PBS; (2) animals receiving a mixture of muFc-Angio and muFc-Endo; (3) animals receiving huKS-muγ2a-muIL2 fusion protein; and (4) animals receiving both the huKS-muγ2a-muIL2 fusion protein and the mixture of muFc-Angio and muFc-Endo. The animals are injected with their respective treatments for five consecutive days.

muFc-Angio and muFc-Endo are administered to tumor-bearing mice at a dose of 5 mg/kg. Animals receiving the huKS-muγ2a-muIL2 fusion protein are treated with daily doses of 10 μg of huKS-muγ2a-muIL2 for a total of 5 days. The growth of the tumors is monitored by volumetric measurements.

It is contemplated that treatment with the mixture of muFc-Angio and muFc-Endo alone will be effective at shrinking LLC tumors over a treatment period of two weeks. It is contemplated, however, that this treatment will not be as effective over a short period of, for example, five days. It is also contemplated that animals treated with the huKS-muIL2 fusion protein alone will exhibit minimal anti-tumor activity. It is contemplated, however, that the group receiving both the huKS-muγ2a-muIL2 fusion protein and a mixture of muFc-Angio and

- 30 -

muFc-Endo will show significantly reduced tumor growth compared to either the muFc-Angio and muFc-Endo group, or the huKS-muγ2a-muIL2 fusion protein group.

**Example 5 Combination Therapy Of Antibody-cytokine Fusion Protein And Endostatin.**

5 Mouse CT26 carcinoma cells expressing human EpCAM were injected subcutaneously in the shaved backs of BALB/c mice ( $2 \times 10^6$  cells per injection). When the tumors reached 100-200 mm<sup>3</sup> in size (about 7 to 14 days), the mice were randomized into four groups, 4 mice per group. Group 1 received intravenous injections of 0.2 mL of PBS daily. Group 2 received intravenous injections of muFc-muEndostatin (320 μg/mouse) in PBS daily. Group 3 received intravenous injections of huKS-huγ4-huIL2 fusion protein (10 μg/mouse) in PBS daily for 5 days only. Group 10 4 received intravenous injections of a combination of huKS-huγ4-huIL2 (10 μg/mouse) and muFc-muEndo (320 μg/mouse) in PBS daily for 5 days, and thereafter daily injections of muFc-muEndo (320 μg/mouse) in PBS. Tumor volumes were measured as described in Example 3.

The results are summarized in FIG. 4. Mice treated with PBS are represented by closed diamonds, mice treated with the muFc-muEndo fusion protein are represented by closed squares, 15 mice treated with a huKS-huγ4-huIL2 fusion protein are represented by closed diamonds, and mice treated with a combination of the muFc-muEndo fusion protein and the huKS-huγ4-huIL2 fusion protein are represented by crosses.

FIG. 4 shows that the combination of the antibody-cytokine fusion protein and the anti-angiogenic protein muFc-muEndo was superior to either agent by itself. After treatment for 19 20 days, the T/C ratio (average size of tumors in the treatment group/average size of tumors in the control group) for the combination therapy of huKS-huγ4-huIL2 and muFc-muEndo was 0.25, which was a significant improvement over the T/C of 0.31 for huKS-huγ4-huIL2 and 0.42 for muFc-muEndo.

**Example 6 Combination Therapy of Antibody-cytokine Fusion Protein and indomethacin.** 25

Female C57B1/6 mice were injected subcutaneously in the mid-back with LLC-Ep cells ( $2 \times 10^6$  cells per injection). When the tumors reached 600-1200 mm<sup>3</sup>, the mice were sacrificed. The skin overlying the tumor was cleaned with betadine and ethanol, the tumors excised and necrotic tissue discarded. A suspension of tumor cells in phosphate buffered saline was prepared by



- 31 -

passing viable tumor tissue through a sieve and then through a series of sequentially smaller hypothermic needles of 22- to 30- gauge. The cells were adjusted to a concentration of  $1 \times 10^7$  cells/mL and placed on ice. C57BL/6 mice then were injected with 0.1 mL of the freshly resuspended cells ( $1 \times 10^6$  cells/mouse) in the proximal midline of the subcutaneous dorsa.

- 5           When the tumors reached 100-200 mm<sup>3</sup> in size (about 7 to 14 days), the mice were randomized into four groups, 5 mice per group. Group 1 received intravenous injections of 0.2 mL of PBS daily. Group 2 received 5 daily intravenous injections of huKS-huyl-huIL2 (25 µg/mouse) in PBS. Group 3 received indomethacin orally in drinking water (20 µg/mL, or about 60-70 µg of indomethacin consumed daily per mouse) throughout the treatment period. Group 4  
10 received 5 daily intravenous injections of huKS-huyl-huIL2(25 µg/mouse), and indomethacin orally in drinking water (20 µg/mL) throughout the treatment period. Tumor volumes were measured as described in Example 3.

- The results are presented in FIG. 5. Mice treated with PBS are represented by closed diamonds, mice treated with the huKS-huyl-huIL2 fusion protein are represented by closed  
15 squares, mice treated with indomethacin are represented by closed triangles, and mice treated with a combination of the huKS-huyl-huIL2 fusion protein and indomethacin are represented by crosses.

- FIG. 5 shows that the combination of an antibody-cytokine fusion protein and the anti-angiogenic chemical compound indomethacin was superior to either agent by itself. After  
20 treatment for 22 days, the T/C ratio for the combination therapy of huKS-huyl-huIL2 and indomethacin was 0.40, which was a significant improvement over the T/C of 0.61 for huKS-huyl-huIL2 and 0.60 for indomethacin.

- 32 -

*Equivalents*

The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting on the invention described herein. Scope of the  
5 invention is thus indicated by the appended claims rather than by the foregoing description, and all changes which come within the meaning and range of equivalency of the claims are intended to be embraced therein.

*Incorporation by Reference*

Each of the patent documents and scientific publications disclosed hereinabove is  
10 incorporated herein by reference.

- 33 -

What is claimed is:

- 1 1. A method of inducing a cytotoxic immune response against a preselected cell-type in a  
2 mammal, the method comprising:  
3 administering to the mammal (i) an immunoconjugate comprising an antibody  
4 binding site capable of binding the preselected cell-type and a cytokine capable of inducing  
5 a said immune response against the preselected cell-type, and (ii) an angiogenesis inhibitor  
6 in an amount sufficient to enhance said immune response relative to immunoconjugate  
7 alone.
- 1 2. The method of claim 1, wherein the preselected cell-type is a cancer cell.
- 1 3. The method of claim 1, wherein the preselected cell-type is a virus-infected cell.
- 1 4. The method of claim 1, wherein the angiogenesis inhibitor is co-administered together  
2 with the immunoconjugate.
- 1 5. The method of claim 1, wherein the angiogenesis inhibitor is administered prior to  
2 administration of the immunoconjugate.
- 1 6. The method of claim 1, wherein the antibody binding site comprises, in an amino-terminal  
2 to carboxy-terminal direction, an immunoglobulin variable region, a CH1 domain, and a  
3 CH2 domain.
- 1 7. The method of claim 6, wherein the antibody binding site further comprises a CH3 domain  
2 attached to the carboxy terminal end of the CH2 domain.
- 1 8. The method of claim 1, wherein the immunoconjugate is a fusion protein comprising, in an  
2 amino-terminal to carboxy-terminal direction, (i) the antibody binding site comprising an  
3 immunoglobulin variable region capable of binding a cell surface antigen on the  
4 preselected cell type, an immunoglobulin CH1 domain, an immunoglobulin CH2 domain,  
5 and (ii) the cytokine.
- 1 9. The method of claim 8, wherein the antibody binding site further comprises a CH3 domain  
2 interposed between the CH2 domain and the cytokine.

- 34 -

- 1 10. The method of claim 1, wherein the cytokine of the immunoconjugate is selected from the  
2 group consisting of a tumor necrosis factor, an interleukin, a colony stimulating factor,  
3 and a lymphokine.
- 1 11. The method of claim 1, wherein the angiogenesis inhibitor is selected from the group  
2 consisting of an endostatin, angiostatin, a peptide having binding affinity for  $\alpha_v\beta_3$  integrin,  
3 an antibody having binding affinity for  $\alpha_v\beta_3$  integrin, a peptide with binding affinity for an  
4 EGF receptor, an antibody having binding affinity for an EGF receptor, a COX-2 inhibitor,  
5 fumagillin, thalidomide, an anti-angiogenic cytokine and a cytokine fusion protein.
- 1 12. A method of inducing a cytotoxic immune response against a cancer cell in a mammal, the  
2 method comprising:  
3 administering to the mammal (i) an immunoconjugate comprising an antibody  
4 binding site capable of binding the cancer cell and a cytokine capable of inducing a said  
5 immune response against the tumor cell, and (ii) an angiogenesis inhibitor selected from  
6 the group consisting of endostatin and angiostatin in an amount sufficient to enhance said  
7 immune response relative to immunoconjugate alone.
- 1 13. The method of claim 12, wherein the angiogenesis inhibitor is co-administered together  
2 with the immunoconjugate.
- 1 14. The method of claim 12, wherein the angiogenesis inhibitor is administered prior to  
2 administration of the immunoconjugate.
- 1 15. The method of claim 12, wherein the antibody binding site comprises, in an amino-  
2 terminal to carboxy-terminal direction, an immunoglobulin variable region, a CH1 domain,  
3 and a CH2 domain.
- 1 16. The method of claim 15, wherein the antibody binding site further comprises a CH3  
2 domain attached to the carboxy terminal end of the CH2 domain.
- 1 17. The method of claim 12, wherein the immunoconjugate is a fusion protein comprising, in  
2 an amino-terminal to carboxy-terminal direction, (i) the antibody binding site comprising  
3 an immunoglobulin variable region capable of binding a cell surface antigen on the  
4 preselected cell type, an immunoglobulin CH1 domain, an immunoglobulin CH2 domain,  
5 and (ii) the cytokine.

- 35 -

- 1 18. The method of claim 17, wherein the antibody binding site further comprises a CH3  
2 domain interposed between the CH2 domain and the cytokine.
- 1 19. The method of claim 12, wherein the cytokine of the immunoconjugate is selected from  
2 the group consisting of a tumor necrosis factor, an interleukin, a colony stimulating factor,  
3 and a lymphokine.
- 1 20. A composition for inducing an immune response against a preselected cell-type in a  
2 mammal, the composition comprising in combination:
- 3 (i) an immunoconjugate comprising an antibody binding site capable of binding the  
4 preselected cell-type and a cytokine capable of inducing an immune response against the  
5 preselected cell-type in the mammal, and
- 6 (ii) an angiogenesis inhibitor in an amount sufficient to enhance said immune  
7 response induced by the immunoconjugate of the combination relative to  
8 immunoconjugate alone.
- 1 21. The composition of claim 20, wherein the antibody binding site comprises in an amino-  
2 terminal to carboxy-terminal direction, an immunoglobulin variable region, a CH1 domain  
3 and a CH2 domain.
- 1 22. The composition of claim 21, wherein the antibody binding site further comprises a CH3  
2 domain attached to the C-terminal end of the CH2 domain.
- 1 23. The composition of claim 20, wherein the immunoconjugate is a fusion protein  
2 comprising, in an amino-terminal to carboxy-terminal direction, (i) the antibody binding  
3 site comprising an immunoglobulin variable region capable of binding a cell surface  
4 antigen on the preselected cell type, an immunoglobulin CH1 domain, an immunoglobulin  
5 CH2 domain, and (ii) the cytokine.
- 1 24. The composition of claim 23, wherein the antibody binding site further comprises a CH3  
2 domain interposed between the CH2 domain and the cytokine.
- 1 25. The composition of claim 20, wherein the cytokine of the immunoconjugate is selected  
2 from the group consisting of a tumor necrosis factor, an interleukin, a colony stimulating  
3 factor, and a lymphokine.

- 36 -

- 1 26. The composition of claim 20, wherein the angiogenesis inhibitor is selected from the group  
2 consisting of an endostatin, angiostatin, a peptide having binding affinity for  $\alpha_v\beta_3$  integrin,  
3 an antibody having binding affinity for  $\alpha_v\beta_3$  integrin, a peptide with binding affinity for an  
4 EGF receptor, an antibody having binding affinity for an EGF receptor, a COX-2 inhibitor,  
5 fumagillin, thalidomide, an anti-angiogenic cytokine and a cytokine fusion protein.
- 1 27. The composition of claim 20, wherein the preselected cell-type is a cancer cell.

1/5




FIG. 1

2/5

FIG. 2A





FIG. 2B





3/5




FIG. 3

4/5




FIG. 4

5/5




FIG. 5

## SEQUENCE LISTING

<110> Gillies, Stephen D

<120> Enhancement of Antibody-Cytokine Fusion Protein  
Mediated Immune Responses by Coadministration with  
Angiogenesis Inhibitor

<130> LEX-004PC

<140>

<141>

<150> 60/081,863

<151> 1998-04-14

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<220>

<221> misc\_feature

<222> (12)..(14)

<223> Translation initiation codon

<400> 1

tctagagcag catggcgccc ccgc 24

<210> 2

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<220>

<221> misc\_feature

<222> (7)..(9)

<223> Translation termination anticodon

<400> 2

ctcgagttat gcattgagtt ccct 24

<210> 3

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 3

ccgtctcctc agccaaaaca acagcccat cggtc 35

<210> 4

- 2 -

<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 4  
ggggctgttg ttttggtga ggagacggtg actgacg 37

<210> 5  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 5  
cttaagccag atccagttgg tgcag 25

<210> 6  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 6  
cccggggtcc gggagaagct cttagtc 27

<210> 7  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 7  
ggccccgggta aagcaccac ttcaagctcc 30

<210> 8  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 8  
ccctcgagtt attgagggtg tgttg 25

<210> 9  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 9

- 3 -

ggaaataaaa cgggctgatg ctgcaccaac tg 32

<210> 10  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 10  
gcagcatcag cccgttttat ttccagcttg gtcc 34

<210> 11  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 11  
cttaagcgag atcgtgctga cccag 25

<210> 12  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 12  
ctcgagctaa cactcattcc tgttgaagc 29

<210> 13  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 13  
ccccgggtag ggtcattcca gtctctgg 28

<210> 14  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:PCR Primer

<400> 14  
ctcgagtcag gcggagctca gatagc 26

<210> 15  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>

- 4 -

&lt;223&gt; Description of Artificial Sequence:PCR Primer

&lt;400&gt; 15

tctagaccat gtgtcctcag aagctaac

28

&lt;210&gt; 16

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PCR primer

&lt;400&gt; 16

ctcgagctag gatcggaacc tgcag

25

&lt;210&gt; 17

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PCR primer

&lt;400&gt; 17

ccccaagctt catactcatc aggactttc

29

&lt;210&gt; 18

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PCR primer

&lt;400&gt; 18

cccctcgagc tatttgaga aagaggtc

28

&lt;210&gt; 19

&lt;211&gt; 49

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PCR primer

&lt;400&gt; 19

ccccaagctt gtgtatctgt cagaatgtaa gccctcctgt ctctgagca

49

&lt;210&gt; 20

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:PCR primer

&lt;400&gt; 20

cccctcgagc taccctcctg tctctgagca

30